Patient-Safety in Africa 2023

Guest contribution: David Guisselquist, 09.05.2023

Africans get HIV from healthcare

A missing issue at the 2023 Global Summit on Patient Safety

In late February 2023, I attended the 5th Global Ministerial Summit on Patient Safety in Montreux, Switzerland.[1] Because there was so much good mixed with bad, it's taken me more than a month to assemble these thoughts. Good: I believe that many people at the conference were fully committed to what they professed — better health, safer healthcare. Bad: I found few attendees — patient safety experts and advocates — aware of the decadeslong and continuing failure to address nosocomial (from health care) HIV infections in sub-Sahara Africa.



9a5th Global Ministerial Summit on Patient Safety in Montreux, Switzerland.
ni
za
ti
on
(W
H0
)
he
lp
s
to

gu id е th es е su mm it s. WH 0′ S at te nt io n to ра ti en t sa fe ty bu il ds on а 20 02 Wo rl d He al th As se mb ly re S0 lu ti on (W HA 55 .1 8) [2] ur gi ng me mb er st at es "t 0 ра у th е cl 0 S es t ро SS ib le at te nt io

n to ...р at ie nt sa fe ty ,, an d te 11 in g WH 0 to ad vi se an d he lp th em to do S0 . WH 0 ha S be en sa yi ng

al			
l			
th			
е			
ri			
gh			
t			
th			
in			
gs			
Fo			
r			
ex			
am			
pl			
е,			
а			
20			
19			
Wo			
rl			
d			
Не			
al			
th			
As			
se			
mo			
ty re			
re			
50			
tu ti			
00			
(W			
НА			
72			
.6			
) [
31			

di re ct S WH 0 "t 0 pr ٥v id е te ch ni са ι su рр or t to Ме mb er St at es , es ре ci al ly lo W an d mi dd le

in C0 me CO un tr ie s" to he lp th em "a SS es s, me as ur е an d im pr ov е ра ti en t sa fe ty ..." Re sp on di ng

to th at re S0 lu ti on , WH 0′ S Gl ob al Ра ti en t Sa fe ty Ac ti on Ρl an 20 21 -2 03 0 pr ор 0S es go ve rn me nt

S (р 13 in [4]) to "Е ns ur е а со ns ta nt fl OW of in fo rm at io n an d kn OW le dg е to dr iv e... im pr ٥v em en ts in th sa fe ty of ca re "

Background: investigations outside sub-Saharan Africa

If WHO staff and associated experts were serious about the plans and responsibilities laid out in these documents, they would be aware of unexplained HIV infections in Africa, and would recommend governments to investigate – the appropriate and proven strategy to protect patients. As demonstrated around the world, investigations of unexplained HIV infections: first identify the likely source clinics, next look for more victims by inviting others attending those clinics to come for tests; and from knowing who was infected find and fix dangerous procedures.

Beginning in 1986, investigations around the world – except in sub-Saharan Africa – have uncovered dozens of small to large outbreaks from medical procedures. In 11 countries in Asia, Europe, North Africa, and the Americas investigations uncovered 12 outbreaks with more than 100 to an estimated 100,000 infections.[5]

For example, in early 2019 a private doctor in Ratodero, Pakistan, sent a persistently sick 1-year old girl for an HIV test; she was HIV-positive, but her mother was not. Over the next several months, he found more infected children with HIV-negative mothers. After the media reported these infections in late April 2019, government arranged for widespread HIV tests on demand.[6] As of early 2023, government's investigation had found more than 2,800 children along with hundreds of adults to be HIV-positive in and near Ratodero.[7] How did it happen? In June 2019, UNAIDS' regional director reported: "Unsafe injection practices including reuse of syringes

and IV [intravenous] drips, both by the doctors as well as quacks [unlicensed healthcare providers] in addition to poor infection control have emerged as the leading causes of HIV outbreak in Ratodero…"[8] According to Fatima Mir, a member of the investigating team, "To call [infection control] abysmal doesn't even do justice to how bad it is… [T]here was a constant contamination of needles that were reused and reused and reused."[9]

Investigations work: as of 2019, WHO estimated adult HIV prevalence of 0.2% or less in 9 of the 11 countries with large investigated outbreaks.[5]

Background: unexplained HIV infections in sub-Saharan Africa

From the mid-1980s and continuing, surveys and studies across sub-Saharan Africa have reported bunches of unexplained HIV infections. For example, during 2006-18, in national surveys in countries with at least 5% of adults HIV-positive, the percentage of self-declared virgins aged 15-24 years who tested HIV-positive ranged as high as 6.3% for men and 5.0% for women (Figure 4.1 in [5]). In seven national surveys during 2006-17 that tested mothers and children for HIV, the percentage of HIV-positive children with mothers testing HIV-negative ranged from 6% to 33% (Figure 4.2 in [5]); 33% was for children aged 6-23 months in Mozambique in 2015.

All along, studies in Africa looking for HIV infections and risks have reported bunches of unexplained infections. For example, a 2011-12 study among young women in Mpumalanga Province, South Africa, tested 2,533 high school women aged 13-20 years; 81 were HIV-positive, including 38 who reported never having vaginal or anal sex.[10] The study then followed and retested the women for 1-6 years during which time 190 got HIV, including 44 who reported no lifetime sex.[11]

Studies and surveys are only the tip of the iceberg of unexplained infections. With huge increases in HIV testing following the UN's 2016 target for 90% of HIV-positive people to know their status by 2020,[12] hundreds of thousands of Africans now know they are infected despite no sexual or mother-to-child risks. In Eastern and Southern Africa, the region with the worst epidemics, 90% of those infected knew their status as of 2021, including 92%-94% in Botswana, eSwatini, Lesotho, and South Africa, the four countries with the worst epidemics, with 18.3%-27.9% adult HIV prevalence (WHO estimates for 2021[13]). Even so, not all unexplained infections have been recognized, because many HIV-positive people with no sex or mother-to-child risks have seen no need to test.

Not protecting patients in sub-Sahara Africa from nosocomial HIV

As demonstrated in countries throughout the world, both the general public and health experts have recognized unexplained infections as evidence patients are at risk unless and until investigations find and fix their source. In other words, government decisions to investigate have nothing to do with the percentages of HIV infections in a country or community from sex or bloodborne risks. Patient risk is the issue.

However, despite World Health Assembly resolutions charging governments and WHO to work together to protect patient safety, WHO and governments have ignored the threat to patient safety represented by unexplained infections in sub-Saharan Africa. Continuing failure to investigate – or even to talk about and advocate investigations – makes a mockery of celebrated, meaningful, and articulated commitments to protect patients.

Letting go of the tiger's tail?

WHO and associated public health experts have a tiger by the tail. They grabbed the tail in the 1980s by not investigating recognized unexplained infections, and not warning Africans about unaddressed risks to get HIV from healthcare. At that time, Africa's HIV epidemics were not so large – as of 1988, WHO estimated 2.5 million infected in sub-Sahara Africa, about 0.6% of the population. But the tiger grew huge – as of end-2021, WHO estimates 28 million Africans had already died of AIDS and 25 million were living with HIV.

Recent HIV sequencing studies provide strong evidence that bloodborne, not sexual, transmission drives Africa's HIV epidemics. A 2022 review[14] of sequencing studies in Africa found five that sequenced HIV collected from large percentages of infected adults in geographically defined communities and then looked for similar viruses showing linked infections. These five studies identified sex partners with similar viruses to explain only 0.3% to 7.5% of HIV-positive adults with sequenced HIV in each study. Moreover, across eight studies that reported the sex of persons with similar infections, a median of 53% of sequence pairs linked persons of the same sex; in other words, a woman was equally likely to get HIV from a woman as from a man, which suggests that most HIV transmission had nothing to do with sex. If, as these studies suggest, bloodborne transmission drives Africa's epidemics, investigations that find and stop bloodborne risks could not only protect patients but also mark the beginning of the end of Africa's HIV epidemics.

Will WHO let go of the tiger's tail? Based on what I saw at the Summit on Patient Safety in Montreux in February 2023, I am not hopeful WHO and associated experts will reconsider and revise their deadly silence about investigations as the standard response to unexplained infections.

After Montreux, here's what I expect: within not more than several years, the African general public, made aware of unexplained infections by expanded HIV testing, will push their governments to investigate, as happened in Ratodero in 2019. That will take care of the issue, at least as far as Africans' patient safety and HIV epidemics are concerned. But once that's done, how will health professionals who ignored this disaster for decades – some misdirected the response, most trustingly accepted misdirection – respond when investigations discover nosocomial HIV outbreaks in Africa? Findings that are good for African's health may also be good also for health professionals by helping them to come in from the cold; ignoring or denying commitments and ideals is not a good way to live.

References

- Federal Office of Public Health, Switzerland. Patient Safety: 5th Global Ministerial Summit 2023. Available at: https://pss2023.ch/
- 2. World Health Assembly (WHA). WHA55: Quality of care: patient safety. Geneva: WHA, 2002. Available at: https://apps.who.int/gb/archive/pdf files/WHA55/ewha5518.pdf,
- 3. WHA. WHA72.6: Global action on patient safety. Geneva: WHA, 2019. Available at:

```
https://apps.who.int/gb/ebwha/pdf_files/WHA72/A72_R6-en.pdf
```

4. World Health Organization (WHO). Global patient safety action plan 2021-2030: toward eliminating avoidable harm in health care. Geneva: WHO, 2021. Available at: https://www.who.int/teams/integrated-health-services/patient-safety/poli
cy/global-patient-safety-action-

plan#:~:text=The%20action%20plan%20aims%20to,years%20(2021%E2%80%932030)

- 5. Ouyang H. The City Losing Its Children to H.I.V. New York Times Magazine 2 April 2021. Available at: https://www.nytimes.com/2021/03/31/magazine/pakistanhiv.html (accessed 8 April 2021).
- 6. Bhatti MW. Dozens getting HIV positive on weekly basis in four Sindh talukas. Geo News [internet] 7 March 2023. Available at: https://www.google.com/search?q=ratodero+hiv&tbm=nws&sxsrf=AJOqlzWmi E2vmT8d8hYhDlQ9NObo0DKAPg:1678603074649&source=lnt&tbs=qdr:w&sa=X&ved=2a hUKEwjdze7f49X9AhWLk4sKHZikDmUQpwV6BAgBEBc&biw=1366&bih=600&dpr=1 (acces sed 12 March 2023).
- 7. Bhatti MW. UNAIDS delegation briefs Sindh Governor of causes of HIV outbreak in Ratodero, Larkana. *TheNews* 14 June 2019.
- 8. Green A. HIV epidemic in children in Pakistan raises concern. *Lancet* 2019; 393: 2288.
- 9. Gisselquist D. Stopping Bloodborne HIV: investigating unexplained infections. London: Adonis & Abbey, 2021. Available at: https://sites.google.com/site/davidgisselquist/stoppingbloodbornehiv (accessed 5 May 2023).
- Pettifor A, MacPhail C, Selin A, et al. HPTN 068: a randomized control trial of a conditional cash transfer to reduce HIV infection in young women in South Africa – study design and baseline results. AIDS Behav 2016; 9: 1863-1882.
- 11. Stoner MCD, Nguyen N, Kilburn K, et al. Age-disparate partnerships and incident HIV infection in adolescent girls and young women in rural South Africa. AIDS 2019; 33: 83-91.
- 12. United Nations (UN). General Assembly reslution 70/266: Political Declaration on HIV and AIDS: On the Fast Track to Accelerating the Fight against HIV and to Ending the AIDS Epidemic by 2030. New York: UN, 2016. Available at:

https://www.unaids.org/sites/default/files/media_asset/2016-political-de claration-HIV-AIDS_en.pdf (accessed 8 May 2023).

- WHO. HIV estimates with uncertainty bounds, 1990-2021. Geneva: WHO, 2022.
- 14. Gisselquist D. Recognizing and stopping blood-borne HIV transmission in Africa. Social Science Research Network (internet), 2022. Available at: https://papers.ssrn.com/sol3/papers.cfm?abstract_id=4174723 (accessed 6 May 2023).

Related Articles

- \bullet AIDS and Patient-Safety in Africa -
- AIDS: Why Africa? -
- Ebola: Why? -
- Unsafe injections 1985-1990 -
- \bullet Management of errors & development cooperation -
- \bullet The development of the human ecosystem -
- Nosocomial viral infections -
- Hepatitis C eradication -
- Aus Fehlern Lernen (Learning from mistakes?) -